



## Similarities and Differences between Flu and COVID-19



Getting a flu vaccine during 2020-2021 is more important than ever because of the ongoing COVID-19 pandemic.

### What is the difference between Influenza (Flu) and COVID-19?

Influenza (Flu) and COVID-19 are both contagious respiratory illnesses, but they are caused by different viruses. COVID-19 is caused by infection with a new coronavirus (called SARS-CoV-2), and flu is caused by infection with [influenza viruses](#).

COVID-19 seems to spread more easily than flu and causes more serious illnesses in some people. It can also take longer before people show symptoms and people can be contagious for longer. More information about differences between flu and COVID-19 is available in the different sections below.

Because some of the symptoms of flu and COVID-19 are similar, it may be hard to tell the difference between them based on symptoms alone, and [testing](#) may be needed to help confirm a diagnosis.

While more is learned every day about COVID-19 and the virus that causes it, there is still a lot that is unknown. This page compares COVID-19 and flu, given the best available information to date.

Learn more about how to protect yourself and others from COVID-19.



Learn more about how to protect yourself and others from flu this season.



### Signs and Symptoms

#### Similarities:

Both **COVID-19** and **flu** can have varying degrees of signs and symptoms, ranging from no symptoms (asymptomatic) to severe symptoms. Common symptoms that COVID-19 and flu share include:

- Fever or feeling feverish/chills
- Cough
- Shortness of breath or difficulty breathing
- Fatigue (tiredness)
- Sore throat
- Runny or stuffy nose
- Muscle pain or body aches
- Headache



- Some people may have vomiting and diarrhea, though this is more common in children than adults

## Differences:

### Flu

Flu viruses can cause mild to severe illness, including common signs and symptoms listed above.

[Flu Symptoms](#)

### COVID-19

COVID-19 seems to cause more serious illnesses in some people. Other signs and symptoms of COVID-19, different from flu, may include change in or loss of taste or smell.

[COVID-19 Symptoms](#)

## How long symptoms appear after exposure and infection

### Similarities:

For both COVID-19 and flu, 1 or more days can pass between a person becoming infected and when he or she starts to experience illness symptoms.

### Differences:

If a person has COVID-19, it could take them longer to develop symptoms than if they had flu.

### Flu

Typically, a person develops symptoms anywhere from **1 to 4 days after infection**.

[Flu Symptoms](#)

### COVID-19

Typically, a person develops symptoms **5 days after being infected**, but symptoms can appear **as early as 2 days after infection** or **as late as 14 days after infection**, and the time range can vary.

[COVID-19 Symptoms](#)



## How long someone can spread the virus

### Similarities:

For both COVID-19 and flu, it's possible to spread the virus for at least 1 day before experiencing any symptoms.

### Differences:

If a person has COVID-19, they may be contagious for a longer period of time than if they had flu.

### Flu

Most people with flu are contagious for about 1 day before they show symptoms.

Older children and adults with flu appear to be most contagious during the initial 3-4 days



of their illness but many remain contagious for about 7 days.

Infants and people with weakened immune systems can be contagious for even longer.

[How Flu Spreads](#)

**COVID-19**

*How long someone can spread the virus that causes COVID-19 is still under investigation.*

It's possible for people to spread the virus for about 2 days before experiencing signs or symptoms and remain contagious for at least 10 days after signs or symptoms first appeared. If someone is asymptomatic or their symptoms go away, it's possible to remain contagious for at least 10 days after testing positive for COVID-19.

[How COVID-19 Spreads](#)

How it Spreads

Similarities:

**Both COVID-19 and flu** can spread from person to person, between people who are in close contact with one another (within about 6 feet). Both are spread mainly by droplets made when people with the illness (COVID-19 or flu) cough, sneeze, or talk. These droplets can land in the mouths or noses of people who are nearby or possibly be inhaled into the lungs.

It may be possible that a person can get infected by physical human contact (for example, shaking hands) or by touching a surface or object that has virus on it and then touching their own mouth, nose, or possibly their eyes.

Both flu virus and the virus that causes COVID-19 may be spread to others by people before they begin showing symptoms; by people with very mild symptoms; and by people who never developed symptoms (asymptomatic).

Differences:

While COVID-19 and flu viruses are thought to spread in similar ways, COVID-19 is more contagious among certain populations and age groups than flu. Also, COVID-19 has been observed to have more superspreading events than flu. This means the virus that causes COVID-19 can quickly and easily spread to a lot of people and result in continuous spreading among people as time progresses.

[How Flu Spreads](#)

[How COVID-19 Spreads](#)



People at High-Risk for Severe Illness

Similarities:

**Both COVID-19 and flu** illness can result in severe illness and complications. Those at highest risk include:

- Older adults
- People with certain underlying medical conditions
- Pregnant people

Differences:

The risk of complications for healthy children is higher for flu compared to COVID-19.



However, infants and children with underlying medical conditions are at increased risk for both flu and COVID-19.

Flu

Young children are at higher risk of severe illness from flu.

[People at High Risk for Flu Complications](#)

COVID-19

School-aged children infected with COVID-19 are at higher risk of [Multisystem Inflammatory Syndrome in Children \(MIS-C\)](#), a rare but severe complication of COVID-19.

[People at Increased Risk of COVID-19 Severe Illness](#)

Complications

Similarities:

Both COVID-19 and flu can result in complications, including:

- Pneumonia
- Respiratory failure
- Acute respiratory distress syndrome (fluid in the lungs)
- Sepsis
- Cardiac injury (for example, heart attacks and stroke)
- Multiple-organ failure (respiratory failure, kidney failure, shock)
- Worsening of chronic medical conditions (involving the lungs, heart, or nervous system or diabetes)
- Inflammation of the heart, brain, or muscle tissues
- Secondary bacterial infections (infections that occur in people who have already been infected with flu or COVID-19)



Differences:

Flu

Most people who get flu will recover in a few days to less than two weeks, but some people will develop [complications](#). Some of these complications are listed above.

[Flu complications](#)

COVID-19

Additional complications associated with COVID-19 can include:

- Blood clots in the veins and arteries of the lungs, heart, legs or brain
- [Multisystem Inflammatory Syndrome in Children \(MIS-C\)](#)

[COVID-19 Emergency warning signs](#)

Approved Treatments

Similarities:

People at high-risk of complications or who have been hospitalized for **COVID-19 or flu**

should receive supportive medical care to help relieve symptoms and complications.

## Differences:

### Flu

Prescription [influenza antiviral drugs](#) are FDA-approved to treat flu.

People who are hospitalized with flu or at high-risk of flu complications with flu symptoms are recommended to be treated with antiviral drugs as soon as possible.

### [Flu Treatment](#)

### COVID-19

The National Institutes of Health (NIH) has developed [guidance on treatment of COVID-19](#) [↗](#), which will be regularly updated as new evidence on treatment options emerges.

While remdesivir is an antiviral agent that is being explored as a treatment for COVID-19 and is available under an Emergency Use Authorization (EUA), there are currently no drugs or other therapeutics approved by the Food and Drug Administration (FDA) to prevent or treat COVID-19. Studies are in progress to learn more.

### [What to Do If You Are Sick with COVID-19](#)



## Vaccine

### Similarities:

Vaccines for **COVID-19** and **flu** must be approved or authorized for emergency use (EUA) by the FDA.

### Differences:

#### Flu

There are multiple FDA-licensed [influenza vaccines](#) produced annually to protect against the 3 or 4 flu viruses that scientists anticipate will circulate each year.

### [Flu Vaccines](#)

### COVID-19

Two [COVID-19 vaccines](#) have been approved for use by the FDA under an EUA. Other vaccines to prevent COVID-19 are still under development.

### [Prevent Getting Sick with COVID-19](#)

